Painkiller Abuse Fuels Rise in Heroin Addiction

Lock Up Your Meds!It could save a life!

For more information on keeping your child safe, visit drugfreenh.org or our website shown on the back of this brochure.



The Surgeon General's
Call to Action says "It's
time to change how
we all think, talk, and
act when it comes to
underage drinking. We
need to stop accepting it
and to start discouraging
it. It's time to help young
people understand that it is
not okay for them to drink
alcohol. The discussion
needs to start long before
youth start thinking about
drinking."

Prescription Drug Misuse and Underage Drinking are serious risk factors for our youth that can have lifelong unintended consequences.

Please talk to your kids — they listen! **Visit drugfreenh.org for information and resources**.

#### SPONSORED BY:








Anonymously dispose of unneeded medications at the Amherst, Hudson, Litchfield, Merrimack, Nashua or Pelham police departments.

To access an online treatment locator in your area please visit www.nhcenterforexcellence.org

# ACKNOWLEDGEMENT & THANK YOU TO:

Vanessa DeZorzi, Graphic Artist, who revised this brochure, Nashua Prevention Coalition, The CoRe Youth Leaders who designed and developed the artwork for this brochure.



# THE PROBLEMS FACING YOUTH AND HOW TO HELP

## **ALCOHOL USE 12-20 year olds**



of New Hampshire 12-20 year olds report drinking alcohol in the past 30 days.

Comparison: 25% Nationally 29% Northeast

Source: 2012 National Survey on Drug Use and Health (NSDUH).

#### PRESCRIPTION DRUG USE

12-17 year olds



of 16-17 year olds in NH have admitting to abusing prescription drugs.

4.6% of NH's 15 year olds admit the same

Source: 2013 Youth Risk Behavior Survey

## NH YOUTH: 1ST TIME **ALCOHOL USE**

When Asked: How old were you when you had your first drink of alcohol other than a few sips?





11.9% 20.9% 21.3%

Responded: Responded: Responded: Responded: 12 or younger

13 or 14

15 or 16

17 or older

Source: 2013 Youth Risk Behavior Survey

## DID YOU KNOW?

Children of parents who talk to their teens regularly about alcohol and other drugs are 42% less likely to use drugs than those who don't. Visit "Talk. They Hear You" at www.underagedrinking.samhsa.gov for age appropriate conversations.

Source: 2013 Youth Risk Behavior Survey

#### **ACCESSIBILITY**

When Asked:

If you wanted to get a prescription drug (such as Oxycontin, Percocet, Vicodin, Codeine, Adderall, Ritalin, or Xanax) without a doctor's prescription, how hard or easy would it be for you to get some?

NH Teens Responded:



said "very easy"

said "sort of easy"

Source: 2013 Youth Risk Behavior Survey

72.4%



Compared to 90.5% nationally, only 72.4% of NH teens think their parents would strongly disapprove of them having one or two drinks of an alcoholic beverage nearly everyday.

Source: 2013 Youth Risk Behavior Survey

### CASES

The following are incidents of two children in NH who accidentally ingested prescription medication in 2014

Age	3	2
Gender	Female	Female
Substance	Insulin	Synthroid/ Vicodin
Action	Observed/ Released	Treated/ Released
Cost	\$1,833.43	\$2,662.76

All info graphics courtesy checkthestatsnh.org

#### **MONITOR:**

Keep track of how many pills you have in the bottle or medication dispenser.

Don't keep alcohol where teens can access it.

Always know how many beers you have in the fridge right now.

Don't allow your teen to have unsupervised parties.

Make sure other parents, grandparents, and adults take the same precautions.

#### SECURE:

Keep prescription drugs and medications in a secure (locked) location.

Lock your liquor cabinet, especially when your child or teen will be unsupervised.

Make sure other parents, grandparents, and adults take the same precautions.

### **DISPOSE:**

Take an inventory of all the prescription drugs in your home. Discard all expired or unused medications.

Discard unused or expired medications by bringing them to local "Take Back Events".

If your local community does not currently support a "Take Back Event", put the unused medication in a plastic bag with used kitty litter, used coffee grinds, or another undesirable substance. Place in the trash.

DO NOT flush medications, unless otherwise specified on the bottle.

Ensure your families privacy. Remove and destroy the labels on the prescription bottle.